

Properly Coloured Cycles and Paths: Results and Open Problems

Gregory Gutin* Eun Jung Kim†

Abstract

In this paper, we consider a number of results and seven conjectures on properly edge-coloured (PC) paths and cycles in edge-coloured multigraphs. We overview some known results and prove new ones. In particular, we consider a family of transformations of an edge-coloured multigraph G into an ordinary graph that allow us to check the existence PC cycles and PC (s, t) -paths in G and, if they exist, to find shortest ones among them. We raise a problem of finding the optimal transformation and consider a possible solution to the problem.

1 Introduction

The class of edge-coloured multigraphs generalize directed graphs. There are several other generalizations of directed graphs such as arc-coloured digraphs, hypertournaments and star hypergraphs, but the class of edge-coloured multigraphs has been given the main attention in graph theory literature because many concepts and results on directed graphs can be extended to edge-coloured multigraphs and there are several important applications of edge-coloured multigraphs. For a more extensive treatment of this topic, see [6, 7].

In this paper we overview some known results on properly coloured (PC) cycles and paths in edge-coloured multigraphs, prove new ones and consider several open problems on the topic. In Section 2 we briefly consider a problem of whether an edge-coloured graph has a PC cycle. In Sections 3 and 4, we offer a useful tool to study edge-coloured multigraphs. In investigating problems on PC subgraphs of edge-coloured multigraphs, it is convenient to transform an edge-coloured graph into an ordinary graph. We suggest a new technique that somewhat automates this transformation. Moreover,

*Department of Computer Science, Royal Holloway, University of London, Egham, Surrey TW20 0EX, UK, [gutinin@cs.rhul.ac.uk](mailto:gutin@cs.rhul.ac.uk)

†Department of Computer Science, Royal Holloway, University of London, Egham, Surrey TW20 0EX, UK, eunjung@cs.rhul.ac.uk

by proving some new results, we illustrate how the proposed technique allows us to obtain more efficient algorithms for PC cycle and PC (s, t) -path problems by reducing the order and size of the transformed graph. We raise a problem of determining the minimum order and size of the transformed graph, and describe the family of graphs that may be the solution to the problem.

In Section 5 we study long PC cycles and paths in arbitrary edge-coloured multigraphs and Section 6 is devoted to longest (mostly Hamilton) PC cycles in edge-coloured complete graphs.

An **m -path-cycle subgraph** F of a multigraph G is a vertex-disjoint union of m paths and a number of cycles in G (some cycles can be of length 2). If $m = 0$, we call F a **cycle subgraph** of G . For a vertex set X of a multigraph G , $G\langle X \rangle$ denotes the subgraph of G induced by X . For a pair s, t of distinct vertices of G , a path between s and t is called an **(s, t) -path**.

We consider **edge-coloured multigraphs**, i.e., undirected multigraphs in which each edge has a colour, but no parallel edges have the same colour. If an edge-coloured multigraph G has c colours, we assume that the colours are $1, 2, \dots, c$ and we call G a **c -edge-coloured** multigraph. We denote the colour of an edge e of an edge-coloured multigraph G by $\chi(e)$. When G has no parallel edges, we call G an **edge-coloured graph**.

Let G be a c -edge-coloured multigraph and let $v \in V(G)$. By $N_i(v)$ we denote the set of neighbours of v adjacent to v by an edge of colour i ; let $d_i(v) = |N_i(v)|$. The **maximum (minimum) monochromatic degree** of $G = (V, E)$ is defined by

$$\begin{aligned} \Delta_{\text{mon}}(G) &= \max\{d_j(v) : v \in V, 1 \leq j \leq c\} \\ \delta_{\text{mon}}(G) &= \min\{d_j(v) : v \in V, 1 \leq j \leq c\}. \end{aligned}$$

Let $\chi(v) = \{i : 1 \leq i \leq c, N_i(v) \neq \emptyset\}$. A path or cycle Q of G is **properly coloured (PC)** if every two adjacent edges of Q are of different colours.

2 Existence of PC Cycles

Since a pair of parallel edges in a c -edge-coloured multigraph ($c \geq 2$) forms a PC cycle, in this section, we consider only c -edge-coloured graphs.

It is easy to see that the problem of checking whether a c -edge-coloured graph has a PC cycle is more general (even for $c = 2$) than the simple problem of verifying whether a digraph contains a directed cycle. Indeed, consider a digraph D and, to obtain a 2-edge-coloured graph G from D , replace each arc xy of D with edges xz_{xy} and $z_{xy}y$ of colours 1 and 2, where z_{xy} is a new vertex ($z_{xy} \neq z_{x'y'}$ provided $xy \neq x'y'$). Observe that G has a PC cycle if and only if D has a directed cycle.

The following theorem by Yeo [19] provides a simple recursive way of checking whether a c -edge-coloured graph has a PC cycle. (For $c = 2$, Theorem 2.1 was first proved by Grossman and Häggkvist [12].)

Theorem 2.1. *Let G be a c -edge-coloured graph, $c \geq 2$, with no PC cycle. Then, G has a vertex $z \in V(G)$ such that no connected component of $G - z$ is joined to z with edges of more than one colour.*

Let us consider the following function introduced by Gutin [13]: $d(n, c)$, the minimum number k such that every c -edge-coloured graph of order n and minimum monochromatic degree at least k has a PC cycle. It was proved in [13] that $d(n, c)$ exists and that

$$d(n, c) \leq \frac{1}{\lfloor c/2 \rfloor} (\log_2 n - \frac{1}{3} \log_2 \log_2 n + \Theta(1)). \quad (1)$$

Abouelaoualim et al. [1] stated a conjecture which implies that $d(n, c) = 1$ for each $c \geq 2$. Using a recursive construction inspired by Theorem 2.1 of c -edge-coloured graphs with minimum monochromatic degree p and without PC cycles, Gutin [13] showed that

$$d(n, c) \geq \frac{1}{c} (\log_c n - \log_c \log_c n) \quad (2)$$

and, thus, the conjecture does not hold. The bounds (1) and (2) imply that $d(n, c) = \Theta(\log_2 n)$ for every fixed $c \geq 2$.

Conjecture 2.2. [13] *There is a function $s(c)$ dependent only on c such that $d(n, c) = s(c) \log_2 n (1 + o(1))$.*

In particular, it would be interesting to determine $s(2)$.

3 P-Gadgets

We consider gadget constructions which generalize some known constructions mentioned below. The P-gadget graphs G^* and G^{**} of an edge-coloured multigraph G described in the next section allow one to transform several problems on properly coloured subgraphs of G into perfect matching problems in G^* or G^{**} .

Let G be an edge-coloured multigraph and let $G' = G - \{x \in V(G) : |\chi(x)| = 1\}$. For each $x \in V(G')$ let G_x be an arbitrary (non-edge-coloured) graph with the following four properties:

P1 $\{x_q : q \in \chi(x)\} \subseteq V(G_x)$;

P2 G_x has a perfect matching;

P3 For each $p \neq q \in \chi(x)$, if the graph $G_x - \{x_p, x_q\}$ is not empty, it has a perfect matching;

P4 For each set $L \subseteq \chi(x)$ with at least 3 elements; if the graph $G_x - \{x_l : l \in L\}$ is not empty, it has no perfect matching.

Each G_x with the properties P1-P4 is called a **P-gadget**. Let us consider the following three P-gadgets; the first two are known in the literature and the third one is new.

1. One P-gadget is due to Szeider [17]:

$$V(G_x) = \{x_i, x'_i : i \in \chi(x)\} \cup \{x''_a, x''_b\} \text{ and}$$

$$E(G_x) = \{x''_a x''_b, x'_i x''_a, x'_i x''_b : i \in \chi(x)\} \cup \{x_i x'_i : i \in \chi(x)\}.$$

We will call this the **SP-gadget**.

2. Another gadget is due to Bang-Jensen and Gutin [4]:

$$V(G_x) = \{x_j : j \in \chi(x)\} \cup \{y_j : j \in \chi(x) \setminus \{m, M\}\},$$

where $m = \min \chi(x)$, $M = \max \chi(x)$, and

$$E(G_x) = \{x_j y_k : j \in \chi(x), k \in \chi(x) \setminus \{m, M\}\} \cup \{x_j x_k : j \neq k \in \chi(x)\}.$$

We will call this the **BJGP-gadget**.

3. The following new gadget is a sort of crossover of the above two and is called the **XP-gadget**:

$$V(G_x) = \{x_j : j \in \chi(x)\} \cup \{y_j : j \in \chi(x) \setminus \{m, M\}\},$$

where m and M are defined above, and

$$E(G_x) = \{x_m x_M\} \cup \{x_j y_j, x_m y_j, x_M y_j : j \in \chi(x) \setminus \{m, M\}\}.$$

It is not difficult to verify that the tree P-gadgets indeed satisfy P1-P4. Let $z = |\chi(x)|$. Observe that the SP-gadget has $2z + 2$ vertices and $3z + 1$ edges, the BJGP-gadget $2z - 2$ vertices and $z(3z - 5)/2$ edges, the XP-gadget $2z - 2$ vertices and $3z - 5$ edges. Thus, the XP-gadget has the minimum number of vertices and edges among the three P-gadgets. It is not difficult to verify that the XP-gadget has the minimum number of vertices and edges among all possible P-gadgets for $z = 2, 3, 4$. Perhaps, this is true for any z .

Conjecture 3.1. *The XP-gadget has the minimum number of vertices and edges among all possible P-gadgets for every $z \geq 2$.*

We will see in the next section why minimizing the numbers of vertices and edges in P-gadgets is important for speeding up some algorithms on edge-coloured multigraphs.

4 P-gadget Graphs

Let G be a c -edge-coloured multigraph and let G_x be a P-gadget for $x \in V(G')$. The graph G^* is defined as follows: $V(G^*) = \cup_{x \in V(G')} V(G_x)$ and $E(G^*) = E_1 \cup E_2$, where $E_1 = \cup_{x \in V(G')} E(G_x)$ and $E_2 = \{y_q z_q : y, z \in V(G'), yz \in E(G), \chi(yz) = q, 1 \leq q \leq c\}$.

Let s, t be a pair of distinct vertices of G and let $H = G - \{s, t\}$. Let G^{**} be constructed from H^* by adding s and t and edges $E_3 = \{sx_i : sx \in E(G), \chi(sx) = i\} \cup \{tx_i : tx \in E(G), \chi(tx) = i\}$.

We will denote the number of vertices and edges in multigraph G , G^* and G^{**} by n, m, n^*, m^*, n^{**} and m^{**} , respectively.

The following result relates perfect matchings of G^* with PC cycle subgraphs of G . PC cycle subgraphs are important in several problems on edge-coloured multigraphs (for example, for the PC Hamilton cycle problem), see [6]. Recall that $G' = G - \{x \in V(G) : |\chi(x)| = 1\}$.

Theorem 4.1. *Let G be a connected edge-coloured multigraph such that G' is non-empty. Then G has a PC cycle subgraph with r edges if and only if G^* has a perfect matching with exactly r edges in E_2 .*

Proof: Let M be a perfect matching of G^* with exactly edges

$$x_{p_1}^1 y_{q_1}^1, \dots, x_{p_r}^r y_{q_r}^r$$

in E_2 . For a vertex x of G' , let Q_x be the set of edges in E_2 adjacent to G_x . By P2, each G_x has even number of vertices ($x \in V(G')$) and since M is a perfect matching in G^* , there is even number of edges in Q_x . By P4, Q_x has either no edges or two edges for each $x \in V(G')$. Let X be the set of all vertices $x \in V(G')$ such that $|Q_x| = 2$. Then, by the definition of G^* , $G\langle X \rangle$ contains a PC cycle factor. It remains to observe that $|X| = r$.

Now let F be a PC cycle subgraph of G with r edges. Observe that the edges of F correspond to a set Q of r independent edges of G^* and that either no edges or two edges of Q are adjacent to G_x for each $x \in V(G')$. Now delete the vertices adjacent with Q from each G_x and observe that each remaining non-empty gadget has a perfect matching by P2 and P3. Combining the perfect matchings of the non-empty gadgets with Q , we get a perfect matching of G^* with exactly r edges from E_2 . $\square$

The first part of the next assertion generalizes a result from [4]. The second part is based on an approach which leads to a more efficient algorithm than in [2].

Corollary 4.2. *One can check whether an edge-coloured multigraph G has a PC cycle and, if it does, find a maximum PC cycle subgraph of G in time $O(n^* \cdot (m^* + n^* \log n^*))$. Moreover one can find a shortest PC cycle in G in time $O(n \cdot n^* \cdot (m^* + n^* \log n^*))$.*

Proof: We may assume that G is connected and that G' is not empty. By Theorem 4.1, it is enough to find a perfect matching of G^* containing the maximum number of edges from E_2 . Assign weight 0 (1, respectively) to edges of G^* in E_1 (E_2 , respectively). Now we need to find a maximum weight perfect matching of G^* which can be done in time $O(n^* \cdot (m^* + n^* \log n^*))$ by a matching algorithm in [11].

To find a shortest PC cycle in G , choose a vertex $x \in V(G')$. We will find a shortest PC cycle in G traversing x . By Theorem 4.1, it is enough to find a perfect matching of G^* containing the minimum number of edges from E_2 while containing at least one edge from E_2 so that the corresponding PC cycle in G should be non-trivial. We define the weights on edges of G^* as follows. Assign M , where M is a sufficiently large number, to each edge in E_2 incident with G_x . For all other edges, assign weight 1 (0, respectively) to edges of G^* in E_1 (E_2 , respectively). A maximum weight perfect matching of G^* contains exactly two edges of weight M by P4, and contains the minimum number of edges in E_2 . Finding a maximum weight perfect matching of G^* can be done in time $O(n^* \cdot (m^* + n^* \log n^*))$ and we iterate the process for each $x \in V(G')$. $\square$

The proof of the following result is analogous to the proof of Theorem 4.1.

Theorem 4.3. *Let G be an edge-coloured multigraph and let s, t be a pair of distinct vertices of G . If G^{**} is non-empty, then G has a PC 1-path-cycle subgraph with r edges in which the path is between s and t if and only if G^{**} has a perfect matching with exactly r edges not in E_1 .*

The next assertion generalizes a result from [2].

Corollary 4.4. *Let G be an edge-coloured multigraph. One can check whether there is a PC (s, t) -path in G in time $O(m^{**})$ and if G has one, a shortest PC (s, t) -path can be found in time $O(n^{**} \cdot (m^{**} + n^{**} \log n^{**}))$.*

Proof: Let L be a graph. Given a matching M in L , a path P in L is M -**augmenting** if, for any pair of adjacent edges in P , exactly one of them belongs to M and the first and last edges of P do not belong to M . Consider a perfect matching M of H^* , where $H = G - \{s, t\}$, which is a collection of perfect matchings of G_x for all $x \in V(G')$. The existence of a perfect matching in G_x is guaranteed by P2. Observe that G has a PC (s, t) -path if and only if there is an M -augmenting (s, t) -path P in G^{**} . Since an M -augmenting path P can be found in time $O(m^{**})$ (see [18]), we can find a PC (s, t) -path in G , if one exists, in time $O(m^{**})$.

To find a shortest PC (s, t) -path, we assign each edge in $\bigcup_{x \in V(G')} E(G_x)$ weight 0 and every other edge of G^{**} weight 1. Observe that a minimum

weight perfect matching Q in the weighted G^{**} corresponds to a shortest PC (s, t) -path. Finding a minimum weight perfect matching can be done in time $O(n^{**} \cdot (m^{**} + n^{**} \log n^{**}))$. $\square$

5 Long PC Cycles and Paths

The following interesting result and conjecture were obtained by Abouelaoualim, Das, Fernandez de la Vega, Karpinski, Manoussakis, Martinhon and Saad [1].

Theorem 5.1. [1] *Let G be a c -edge-coloured multigraph G with n vertices and with $\delta_{mon}(G) \geq \lceil \frac{n+1}{2} \rceil$. If $c \geq 3$ or $c = 2$ and n is even, then G has a Hamilton PC cycle. If $c = 2$ and n is odd, then G has a PC cycle of length $n - 1$.*

Conjecture 5.2. [1] *Theorem 5.1 holds if we replace $\delta_{mon}(G) \geq \lceil \frac{n+1}{2} \rceil$ by $\delta_{mon}(G) \geq \lceil \frac{n}{2} \rceil$.*

We cannot replace $\delta_{mon}(G) \geq \lceil \frac{n+1}{2} \rceil$ by $\delta_{mon}(G) \geq \lceil \frac{n-1}{2} \rceil$ due to the following example. Let H_1 and H_2 be c -edge-coloured complete multigraphs (for each pair x, y of vertices and each $i \in \{1, 2, \dots, c\}$ and $j \in \{1, 2\}$, H_j has a edge between x and y of colour i) of order $p + 1$ that have precisely one vertex in common. Clearly, a longest PC cycle in $H_1 \cup H_2$ is of length $p + 1$.

Since the longest PC path problem is $\mathcal{NP}$ -hard, it makes sense to study lower bounds on the length of a longest PC path. The following result was proved by Abouelaoualim et al. [1].

Theorem 5.3. *Let G be a c -edge-coloured graph of order n with $\delta_{mon}(G) = d \geq 1$. Then G has a PC path of length at least $\min\{n - 1, 2\lfloor \frac{c}{2} \rfloor d\}$.*

The authors of [1] raised the following two conjectures.

Conjecture 5.4. *Let G be a c -edge-coloured graph of order n and let $d = \delta_{mon}(G) \geq 1$. Then G has a PC path of length at least $\min\{n - 1, 2cd\}$.*

They also conjectured the following analog of Theorem 5.3 for multigraphs:

Conjecture 5.5. *Let G be a c -edge-coloured multigraph of order n with $\delta_{mon}(G) = d \geq 1$. Then G has a PC path of length at least $\min\{n - 1, 2d\}$.*

6 Longest PC Cycles and Paths in Edge-Coloured Complete Graphs

Let K_n^c denote a c -edge-coloured complete graph with n vertices.

Feng, Giesen, Guo, Gutin, Jensen and Rafiey [10] proved the following:

Theorem 6.1. *A K_n^c ($c \geq 2$) has a PC Hamilton path if and only if K_n^c contains a PC spanning 1-path-cycle subgraph.*

This theorem was first proved by Bang-Jensen and Gutin [4] for the case $c = 2$ and they conjectured that Theorem 6.1 holds for each $c \geq 2$. Theorem 6.1 implies that the maximum order of a PC path in K_n^c equals the maximum order of a PC 1-path-cycle subgraph of K_n^c .

As a result, the problem of finding a longest PC path in K_n^c is polynomial-time solvable for arbitrary $c \geq 2$. To see that a PC 1-path-cycle subgraph of K_n^c can be found in polynomial time, add a pair x, y of new vertices to K_n^c together with all edges needed to have a complete multigraph on $n + 2$ vertices. Let the colour of all edges between x and y and K_n^c be $c + 1$ and let the colour of xy be $c + 2$. Observe that the maximum order of a PC 1-path-cycle subgraph of K_n^c equals the maximum order of a PC cycle subgraph of the $c + 2$ -edge-coloured complete graph described above. It remains to apply Corollary 4.2.

The problem of finding a longest PC cycle K_n^c has not been solved yet for $c \geq 3$ as we will see below. For $c = 2$, Saad [15] found a characterization for longest PC cycles using the following notions. A pair of distinct vertices x, y of G are **colour-connected** if there exist PC (x, y) -paths P and Q such that $\chi(f_P) \neq \chi(f_Q)$ and $\chi(\ell_P) \neq \chi(\ell_Q)$, where f_P and f_Q are the first edges of P and Q , respectively, and ℓ_P and ℓ_Q are the last edges of P and Q , respectively. We say that G is **colour-connected** if every pair of distinct vertices of G is colour-connected. Saad's characterization is as follows.

Theorem 6.2. *The length of a longest PC cycle in a colour-connected K_n^2 is equal to the maximum order of a PC cycle subgraph of K_n^2 .*

Colour-connectivity for K_n^c is an equivalence relation (see [6]). Using Theorem 6.2, Saad [15] showed that the problem of finding a longest PC cycle in K_n^2 is random polynomial. Using a special case of Corollary 4.2, Bang-Jensen and Gutin [5] proved that the problem is, in fact, polynomial-time solvable. Theorem 6.2 implies the following:

Corollary 6.3. *[15] A K_n^2 has a PC Hamilton cycle if and only if K_n^2 is colour-connected and contains a PC cycle factor.*

There is another characterization of K_n^2 with a PC Hamilton cycle due to Bankfalvi and Bankfalvi, see [6]. The straightforward extension of Corollary 6.3 is not true for any $c \geq 3$ [6]. In fact, no characterization of K_n^c with a PC Hamilton cycle is known for any fixed $c \geq 3$ and it is a very interesting problem to obtain such a characterization. Even the following problem by Benkour, Manoussakis, Paschos and Saad [8] is still open.

Problem 6.4. *Determine the complexity of the PC Hamilton cycle problem for c -edge-coloured complete graphs when $c \geq 3$.*

We conjecture that the PC Hamilton cycle problem for c -edge-coloured complete graphs when $c \geq 3$ is polynomial-time solvable.

In absence of characterization of K_n^c with a PC Hamilton cycle, sufficient conditions are interest. Manoussakis, Spyratos, Tuza and Voigt [14] proved the next result.

Proposition 6.5. *If $c \geq \frac{1}{2}(n-1)(n-2) + 2$, then every K_n^c has a PC Hamilton cycle.*

Let $\Delta_{mon}(K_n^c)$ denote the largest monochromatic degree of K_n^c . Bollobás and Erdős [9] posed the following:

Conjecture 6.6. *Every K_n^c with $\Delta_{mon}(K_n^c) \leq \lfloor n/2 \rfloor - 1$ has a PC Hamilton cycle.*

Improving some previous results on this conjecture, Shearer [16] showed that if $7\Delta_{mon}(K_n^c) < n$, then K_n^c has a PC Hamilton cycle. So far, the best asymptotic estimate was obtained by Alon and Gutin [3].

Theorem 6.7. [3] *For every $\epsilon > 0$ there exists an $n_0 = n_0(\epsilon)$ so that for each $n > n_0$, every K_n^c satisfying $\Delta_{mon}(K_n^c) \leq (1 - \frac{1}{\sqrt{2}} - \epsilon)n$ contains a PC Hamilton cycle.*

References

- [1] A. Abouelaoualim, K.Ch. Das, W. Fernandez de la Vega, Y. Manoussakis, C.A. Martinhon, and R. Saad, Cycles and paths in edge-colored graphs with given degrees, Manuscript, 2007.
- [2] A. Abouelaoualim, K.Ch. Das, L. Faria, Y. Manoussakis, C.A. Martinhon, and R. Saad, Paths and Trails in Edge-Colored Graphs, Proc. LATIN'08, Lecture Notes Comput. Sci., 4957 (2008), 723–735.
- [3] N. Alon and G. Gutin, Properly colored Hamilton cycles in edge colored complete graphs. Random Struct. & Alg., 11 (1997), 179–186.
- [4] J. Bang-Jensen and G. Gutin, Alternating cycles and paths in edge-coloured multigraphs: a survey, Discrete Math., 165–166 (1997), 39–60.
- [5] J. Bang-Jensen and G. Gutin, Alternating cycles and trails in 2-edge-coloured multigraphs, Discrete Math., 188 (1998), 61–72.
- [6] J. Bang-Jensen and G. Gutin, Digraphs: Theory, Algorithms and Applications, Springer, London, 2000, Freely available online at www.cs.rhul.ac.uk/books/dbook/.
- [7] J. Bang-Jensen and G. Gutin, Digraphs: Theory, Algorithms and Applications, 2nd Edition, Springer, London, In preparation.
- [8] A. Benkouar, Y. Manoussakis, V. Paschos and R. Saad, On the Complexity of Some Hamiltonian and Eulerian Problems in Edge-colored Complete Graphs, Lect. Notes Computer Sci., 557 (1991), 190–198.

- [9] B. Bollobás and P. Erdős, Alternating Hamiltonian cycles, *Israel J. Math.*, 23 (1976), 126–131.
- [10] J. Feng, H.-E. Giesen, Y. Guo, G. Gutin, T. Jensen and A. Rafiey, Characterization of edge-colored complete graphs with properly colored Hamilton paths, *J. Graph Theory*, 53 (2006), 333–346.
- [11] H.N. Gabow, Data structures for weighted matching and nearest common ancestors with linking, *Proc. SODA'90*, 1990, 434–443.
- [12] J.W. Grossman and R. Häggkvist, Alternating cycles in edge-partitioned graphs, *J. Combin. Theory Ser. B* 34 (1983), 77–81.
- [13] G. Gutin, Note on edge-colored graphs and digraphs without properly colored cycles. *Austral. J. Combin.*, To appear.
- [14] Y. Manoussakis, M. Spyrtos, Zs. Tuza and M. Voigt Minimal colorings for properly colored subgraphs, *Graphs & Combin.*, 12 (1996), 345–360.
- [15] R. Saad, Finding a longest alternating cycle in a 2-edge-coloured complete graph is in RP, *Combin. Prob. & Comput.*, 5 (1996), 297–306.
- [16] J. Shearer, A property of the colored complete graph, *Discrete Math.*, 25 (1979), 175–178.
- [17] S. Szeider, Finding paths in graphs avoiding forbidden transitions, *Discrete Appl. Math.*, 126 (2003), 261–273.
- [18] R.E. Tarjan, *Data structures and network algorithms*, SIAM-Philadelphia, 1983.
- [19] A. Yeo, A note on alternating cycles in edge-coloured graphs, *J. Combin. Theory Ser. B* 69 (1997), 222–225.